



National
Aeronautics and
Space
Administration

NASA Scientific and Technical Document Availability Authorization (DAA)

The DAA approval process applies to all forms of published NASA Scientific and Technical Information (STI), whether disseminated in print or electronically. It is to be initiated by the responsible NASA Project Officer, Technical Monitor, author, or other appropriate NASA official for all presentations, reports, papers, and proceedings that contain NASA STI. Explanations are on the back of this form and are presented in greater detail in NPG 2200.2, "Guidelines for Documentation, Approval, and Dissemination of NASA Scientific and Technical Information."

☐ Original
☐ Modified

I. DOCUMENT/PROJECT IDENTIFICATION

TITLE Gopher Tortoise (<i>Gopherus polyphemus</i>) densities in coastal scrub and slash pine flatwoods.	AUTHOR(S) Breininger, D.R., P.A. Schmalzer, and C.R. Hinkle.
---	---

ORIGINATING NASA ORGANIZATION Kennedy Space Center Biomedical Office	PERFORMING ORGANIZATION (If different) Dynamac Corporation, KSC
--	---

CONTRACT/GRANT/INTERAGENCY/PROJECT NUMBER(S)	DOCUMENT NUMBER(S) Journal of Herpetology 28: 60-65	DOCUMENT DATE 1994
---	---	------------------------------

For presentations, documents, or other STI to be externally published (including through electronic media), enter appropriate information on the intended publication such as name, place, and date of conference, periodical, or journal name, or book title and publisher in the next box. These documents must be routed to the NASA Headquarters or Center Export Control Administrator for approval (see Sections III and VIII).



II. SECURITY CLASSIFICATION

CHECK ONE (One of the five boxes denoting Security Classification must be checked.)

☐ SECRET ☐ SECRET RD ☐ CONFIDENTIAL ☐ CONFIDENTIAL RD ☒ UNCLASSIFIED

III. AVAILABILITY CATEGORY

<input type="checkbox"/> ITAR <input type="checkbox"/> EAR	Export Controlled Document - USML Category Classification Number (ECCN) <i>(Documents marked in this block must have the concurrence/approval of the NASA Headquarters or Center Export Control Administrator (see Section VIII).)</i>
<input type="checkbox"/> TRADE SECRET <input type="checkbox"/> SBIR <input type="checkbox"/> COPYRIGHTED	Confidential Commercial Document (check appropriate box at left and indicate below the appropriate limitation and expiration): <input type="checkbox"/> U.S. Government agencies and U.S. Government agency contractors only <input type="checkbox"/> NASA contractors and U.S. Government only <input type="checkbox"/> U.S. Government agencies only <input type="checkbox"/> NASA personnel and NASA contractors only <input type="checkbox"/> NASA personnel only <input type="checkbox"/> Available only with the approval of issuing office: <input type="checkbox"/> Limited until (date)
<input checked="" type="checkbox"/> PUBLICLY AVAILABLE	Publicly available documents must be unclassified, may not be export controlled, may not contain trade secret or confidential commercial data, and should have cleared any applicable patents application process.

IV. DOCUMENT DISCLOSING AN INVENTION

THIS DOCUMENT MAY BE RELEASED ON (date)	NASA HQ OR CENTER PATENT OR INTELLECTUAL PROPERTY COUNSEL SIGNATURE	DATE
---	--	-------------

V. BLANKET RELEASE (OPTIONAL)

- ☐ All documents issued under the following contract/grant/project number may be processed as checked in Sections II and III.
- ☐ The blanket release authorization granted on (date)
- ☐ is RESCINDED - Future documents must have individual availability authorizations.
- ☐ is MODIFIED - Limitations for all documents processed in the STI system under the blanket release should be changed to conform to blocks as checked in Sections II and III.

VI. AUTHOR/ORIGINATOR VERIFICATION			
I HAVE DETERMINED THAT THIS PUBLICATION:			
<input type="checkbox"/> DOES contain export controlled, confidential commercial information, and/or discloses an invention for which a patent has been applied, and the appropriate limitation is checked in Sections III and/or IV.			
<input checked="" type="checkbox"/> does NOT contain export controlled, confidential commercial information, nor does it disclose an invention for which a patent has been applied, and may be released as indicated above.			
NAME OF AUTHOR/ORIGINATOR	MAIL CODE	SIGNATURE	DATE
David Breininger	DYN-2	<i>[Signature]</i>	4/6/98
VII. PROJECT OFFICER/TECHNICAL MONITOR/DIVISION CHIEF REVIEW			
<input checked="" type="checkbox"/> APPROVED FOR DISTRIBUTION AS MARKED ON REVERSE <input type="checkbox"/> NOT APPROVED			
NAME OF PROJECT OFFICER OR TECH. MONITOR	MAIL CODE	SIGNATURE	DATE
RAYMOND M. WHEELER	JJ-G	<i>[Signature]</i>	4/8/98
VIII. EXPORT CONTROL REVIEW/CONFIRMATION			
<input checked="" type="checkbox"/> Public release is approved <input type="checkbox"/> Export controlled limitation is not applicable			
<input type="checkbox"/> Export controlled limitation is approved <input type="checkbox"/> Export controlled limitation (ITAR/EAR) marked in Section III is assigned to this document:			
USML CATEGORY NUMBER	CCL ECCN NUMBER	HQ OR CENTER EXPORT CONTROL ADMINISTRATOR (as applicable)	DATE
		<i>[Signature]</i>	4/16/98
IX. PROGRAM OFFICE OR DELEGATED AUTHORITY REVIEW			
<input checked="" type="checkbox"/> APPROVED FOR DISTRIBUTION AS MARKED ON REVERSE <input type="checkbox"/> NOT APPROVED CODE M, DAA LETTER, 6/14/94			
NAME OF PROGRAM OFFICE REPRESENTATIVE	MAIL CODE	SIGNATURE	DATE
Audrey Lee Silipo	FF-S2-A	<i>[Signature]</i>	4/14/98
X. DISPOSITION			
THIS FORM, WHEN COMPLETED, IS TO BE SENT TO YOUR CENTER PUBLICATIONS OFFICE			

INSTRUCTIONS FOR COMPLETING THE NASA SCIENTIFIC AND TECHNICAL DOCUMENT AVAILABILITY AUTHORIZATION (DAA) FORM

Purpose. This DAA form is used to prescribe the availability and distribution of all NASA-generated and NASA-funded documents containing scientific and technical information (including those distributed via electronic media such as the World Wide Web and CD-ROM).

Requirements. The author/originator must provide either a suitable summary description (title, abstract, etc.) or a completed copy of the document with this form. This form is initiated by the document author/originator and that individual is responsible for recommending/determining the availability/distribution of the document. The author/originator completes Sections I through III, and VI. The author/originator is also responsible for obtaining information and signature in Section IV to the extent the document discloses an invention for which patent protection has been applied. Subsequent to completion of these sections, the author/originator forwards the document to the appropriate Project Manager/Technical Monitor/Division Chief for further review and approval in Section VII, including a re-review of the planned availability and distribution. Once this approval is obtained, the DAA is forwarded to the NASA Headquarters or Center Export Administrator for completion of Section VIII. It is then forwarded for completion of Section IX to the cognizant NASA Headquarters Program Office or Delegated Authority, who provides final review and approval for release of the document as marked.

When to Use This Form. Documents containing STI and intended for presentation or publication (including via electronic media) must be approved in accordance with the NASA STI Procedures and Guidelines (NPG 2200.2). Documents that are to be published in the NASA STI Report Series must be coordinated with the appropriate NASA Headquarters or Center Scientific and Technical Information Office in accordance with NPG 2200.2. Note that information on the Report Documentation Page (if attached) is not to be entered on the DAA except for title, document date, and contract number.

How to Use this Form. Specific guidelines for each section of this form are detailed below.

I. Document/Project Identification. Provide the information requested. If the document is classified, provide instead the security classification of the title and abstract. (Classified information must not be entered on this form). Include RTOP numbers on the Contract/Grant/Interagency/Project Number(s) line. Provide information on presentations or externally published documents as applicable.

II. Security Classification. Enter the applicable security classification for the document. Documents, if classified, will be available only to appropriately cleared personnel having a "need to know."

III. Availability Category for Unclassified Documents. Check the appropriate category or categories.

Export Controlled Document. If the document is subject to export restrictions (see NPG 2200.2, paragraph 4.5.3), the appropriate restriction must be checked, either International Traffic in Arms Regulations (ITAR) or Export Administration Regulations (EAR), and the appropriate United States Munitions List (USML) category or Commerce Control List (CCL), Export Control Classification Number (ECCN) must be cited.

Confidential Commercial Documents (Documents containing Trade Secrets, SBIR documents, and/or Copyrighted Information). Check the applicable box (see NPG 2200.2 paragraph 4.5.7). When any of these boxes are checked, also indicate the appropriate limitation and expiration in the list to the right of these restrictions. These limitations refer to the user groups authorized to obtain the document. The limitations apply both to the initial distribution of the documents and the handling of requests for the documents. The limitations will appear on and apply to reproduced copies of the document. Documents limited to NASA personnel should not be made available to onsite contractors. If the Available Only With the Approval of Issuing Office limitation is checked, the NASA Center for AeroSpace Information will provide only bibliographic processing and no initial distribution; CASI will refer all document requests to the issuing office.

Publicly Available Document - Unrestricted Distribution. Check this box if the information in the document may be made available to the general public without restrictions (unrestricted domestic and international distribution). If the document is copyrighted (see paragraph 4.5.7.3 in NPG 2200.2), also check the "Copyrighted" box in this section.

IV. Document Disclosing an Invention. This must be completed when the document contains information that discloses an invention (see NPG 2200.2, paragraph 4.5.9). When this box is checked, an additional appropriate availability category must be checked. Use of this category must be approved by NASA Headquarters or Center Patent Counsel or the Intellectual Property Counsel.

V. Blanket Release (Optional). Complete this optional section whenever subsequent documents produced under the contract, grant, or project are to be given the same distribution and/or availability as described in Sections II and III. More than one contract number or RTOP Number can be entered. This section may also be used to rescind or modify an earlier Blanket Release. All blanket releases must be approved by the Program Office or its designee and concurred with by the Office of Management Systems and Facilities.

VI. Author/Originator Verification. Required for all DAA forms.

VII. Project Officer/Technical Monitor/Division Chief Review. The Project Officer/Technical Monitor/Author or Originator Division Chief or above must sign and date the form. The office code and typed name should be entered.

VIII. Export Control Review/Confirmation. This section is to be completed by the authorized NASA Headquarters or Center Export Control Administrator for all documents.

IX. Program Office or Delegated Authority Review. This section is to be completed by the duly authorized official representing the NASA Headquarters Program Office. Any delegation from NASA Headquarters to a NASA Center in accordance with NPG 2200.2 should be entered here.

X. Disposition. For NASA Center use.

Gopher Tortoise (*Gopherus polyphemus*) Densities in Coastal Scrub and Slash Pine Flatwoods in Florida

DAVID R. BREININGER, PAUL A. SCHMALZER, AND C. ROSS HINKLE

The Bionetics Corporation, NASA Biomedical Operations and
Research Office, Mail Code BIO-2, John F. Kennedy Space Center, Florida 32899, USA

ABSTRACT.—Densities of gopher tortoises were compared with habitat characteristics in scrub and in flatwood habitats on the Kennedy Space Center, Florida. Tortoises were distributed widely among habitat types and did not have higher densities in well-drained (oak-palmetto) than in poorly-drained (saw palmetto) habitats. Fall densities of tortoises ranged from a mean of 2.7 individuals/ha in disturbed habitat to 0.0 individuals/ha in saw palmetto habitat. Spring densities of tortoises ranged from a mean of 2.5 individuals/ha in saw palmetto habitat to 0.7 individuals/ha in oak-palmetto habitat. Densities of tortoises were correlated positively with the percent herbaceous cover, an indicator of food resources. Plots were divided into three burn classes; these were areas burned within three years, burned four to seven years, and unburned for more than seven years prior to the study. Relationships between densities of tortoises and time-since-fire classes were inconsistent.

Most studies of the gopher tortoise (*Gopherus polyphemus*) have been in well-drained longleaf pine (*Pinus palustris*)-turkey oak (*Quercus laevis*) communities known as sandhill, or in pine plantations that were once sandhill. Habitat requirements of tortoises are well-drained, loose soil (Landers and Speake, 1980; McRae et al., 1981; Auffenberg and Franz, 1982; Diemer, 1986). Most tortoises on the Kennedy Space Center, Florida (KSC), occupy scrub and pine flatwood habitats (Breininger et al., 1991a). Little is published on tortoise densities in these habitats. Our objectives were to compare densities of tortoises in scrub and pine flatwood habitats and to investigate the influence of time since the last fire on densities of tortoises.

MATERIALS AND METHODS

The KSC is the largest protected barrier island complex on the east coast of Florida. It has both temperate and subtropical plant and animal associations. Topographic relief on KSC ranges from sea level to 3 m. Scrub and pine flatwoods (Abrahamson and Hartnett, 1990; Myers, 1990; Schmalzer and Hinkle, 1992b) are the major upland habitat types and are interspersed among seasonally and permanently flooded swale marshes. The understory of pine flatwoods on KSC essentially is identical to scrub; pine flatwoods have a sparse canopy of slash pines (*P. elliotii*). We do not distinguish scrub and pine flatwoods hereafter. Scrub on KSC has a dense shrub layer dominated by myrtle oak (*Quercus myrtifolia*) and sand live oak (*Q. geminata*) on drier sites, and by saw palmetto (*Serenoa repens*) on wetter sites (Schmalzer and Hinkle, 1992a,

b). Nearly half the scrub is predominated by saw palmetto, and the remainder is predominated by a mixture of scrub oaks and saw palmetto (Breininger et al., 1991b). Oak-palmetto habitat on KSC often is referred to as oak scrub (Myers 1990) or scrubby flatwoods (Abrahamson and Hartnett, 1990). Saw palmetto habitat is sometimes referred to as mesic flatwoods (Abrahamson and Hartnett, 1990). Fire has little influence on species composition because oaks, palmettos, and ericaceous shrubs resprout from rhizomes (Schmalzer and Hinkle, 1992a, b). The dominant herbaceous plant is wiregrass (*Aristida stricta*). Sites that were historically scrub and were cleared ≥ 20 yr before the study were termed "disturbed habitat." These were revegetated by scrub species but remain different from scrub in both the relative abundance of scrub plants and structure (Breininger and Schmalzer, 1990).

We established 112 30 \times 50 m plots in a stratified random design throughout scrub and flatwoods on KSC (Breininger et al. 1991a). This plot size was the largest size that allowed sampling within homogeneous vegetation. Sixteen plots were established in disturbed habitat, and thirty-two plots were established in each of three burn classes of oak-palmetto and saw palmetto habitat. Burn classes included areas burned within three years, burned four to seven years, and unburned for more than seven years prior to the study. We identified all tortoise burrows within each plot in early fall (September–October 1985) and spring (April–May, 1986). Burrows were classified as active, inactive, or abandoned. Active burrows had recent plastral slide

TABLE 1. Mean (± 1 SD) of broad habitat features distinguishing oak-palmetto, saw palmetto, and disturbed habitats on John F. Kennedy Space Center, Florida.

	Habitat types		
	Disturbed (N = 15)	Oak-palmetto (N = 48)	Palmetto (N = 40)
Total shrub cover (%)	63 (29)	98 (3)	96 (8)
Scrub oak cover (%)	28 (29)	57 (24)	14 (9)
Saw palmetto cover (%)	7 (14)	44 (23)	55 (23)
Depth to water table (cm)	84 (39)	84 (38)	55 (24)

marks and footprints (Auffenberg and Franz, 1982). Inactive burrows were recently maintained but lacked fresh sign of use. Abandoned burrows were filled partly with litter or partially caved in. We used a stick method and bucket traps in fall, and a camera system in spring, to determine if adult and subadult burrows were occupied by tortoises (Breininger et al., 1991a). Occupancy of only active burrows was determined in fall; occupancy of both active and inactive burrows was determined in spring. Only one of 44 inactive burrows was occupied by a tortoise; thus, inactive burrows contributed little to densities of tortoises within plots in spring (Breininger et al., 1991a). Burrows ≥ 12 cm wide at the entrance were classified as subadult or adult burrows; burrows < 12 cm wide at the entrance were classified as hatchling or juvenile burrows (Diemer, 1992a). Active juvenile and hatchling burrows were recorded for each plot only during the spring of 1986.

Vegetation measurements were taken at each plot during summer (June–September 1985) using a point-intercept sampling technique (Mueller-Dombois and Ellenberg, 1974). Plots were sampled using a grid of 20 points spaced 5 m apart on four parallel lines across the plot. Lines were spaced 10 m apart. Species of trees, shrubs, herbs, and shrub height were recorded at each point. Slash pine was the only canopy tree present; all other woody plants were shrubs. Water table measurements were taken by coring in fall (September–October 1985) at all plots. Among oak-palmetto and saw palmetto plots, we noted small (< 2 m wide) mechanical disturbances (clearing), which we referred to as mild disturbance.

Of the original 112 plots, eight plots were burned during the study and one was cleared; these were excluded from analyses. Densities of tortoises were calculated for each plot, separately for each season, by summing the number of tortoises occupying the burrows in a plot and dividing by the area of the plot.

The percent shrub cover, scrub oak cover, saw

palmetto cover, herbaceous cover, and non-*Aristida* herbaceous cover were variables determined for each plot by summing the number of points where the variable was present, dividing by the total of 20 grid points, and multiplying by 100. The calculation of each was independent of others. Non-*Aristida* cover includes ground cover by herbs other than wiregrass. A herbaceous cover variable that excluded wiregrass was selected because other herbs may be better predictors of quality food sources (Garner and Landers, 1981; Macdonald and Mushinsky, 1988). The mean shrub height was determined for each plot by summing the height measurements for each point and dividing by 20.

Scrub plots, which did not include plots in disturbed habitat, were classified as an oak-palmetto habitat if they had $\geq 30\%$ oak cover or as a saw palmetto habitat if they had $< 30\%$ oak cover based on field data. Therefore, we analyzed a total of three habitat types: oak-palmetto, saw palmetto, and disturbed. Mean values of habitat features (total shrub cover, scrub oak cover, saw palmetto cover, and distance to water table) were calculated for the three habitat types because these change little with fire history in the burn classes considered here (Schmalzer and Hinkle, 1992a, b). Mean values for habitat variables influenced by fire history (mean shrub height, herbaceous cover, non-*Aristida* herbaceous cover) were calculated for disturbed habitat and for each fire class within oak-palmetto and saw palmetto habitats. Densities of tortoises also were calculated for disturbed habitats and for each fire class within oak-palmetto and saw palmetto habitats. Tortoise densities were not normally distributed in either season (Kolmogorov-Smirnov test [fall]: $z = 5.07$, $P < 0.001$, $N = 103$; [spring] $z = 4.86$, $P < 0.001$, $N = 103$) (SPSS Inc., 1988), preventing the use of parametric statistics.

We calculated Spearman rank correlations among fall and spring tortoise densities and habitat variables by season using data from all plots. The presence or absence of mild distur-

TABLE 2. Mean (± 1 SD) of habitat characteristics and of gopher tortoise densities (individuals/ha) for disturbed habitat and time-since-fire classes (0-3, 4-7, >7 years) within oak-palmetto and saw palmetto habitats.

	Disturbed (N = 15)	Oak-Palmetto				Saw Palmetto			
		0-3 (N = 10)	4-7 (N = 18)	>7 (N = 20)	0-3 (N = 19)	4-7 (N = 9)	>7 (N = 12)		
Habitat characteristics									
Mean shrub height (cm)	130 (76)	80 (22)	92 (15)	156 (66)	79 (24)	106 (11)	137 (27)		
Herbaceous cover (%)	76 (23)	62 (27)	38 (29)	24 (23)	66 (19)	64 (18)	45 (30)		
Non-Aristida herbaceous cover (%)	71 (26)	34 (30)	9 (19)	4 (6)	30 (22)	19 (18)	28 (26)		
Gopher tortoise density									
Fall	2.7 (4.9)	1.3 (2.8)	0.4 (1.6)	0.3 (1.5)	2.1 (3.9)	0.0 (0)	2.2 (4.3)		
Spring	2.2 (4.8)	0.7 (2.1)	1.1 (2.5)	1.3 (2.8)	2.5 (4.0)	1.5 (3.0)	1.1 (2.6)		

bance was treated as a categorical variable to investigate its influence on density of tortoises in oak-palmetto and saw palmetto habitats.

RESULTS

Disturbed habitat did not have as extensive shrub cover as oak-palmetto and saw palmetto habitat (Table 1). Mean depth to the water table was <1 m below the surface in all habitat types. Disturbed habitat had the most herbaceous cover (Table 2). Most of the herbaceous cover was wiregrass in oak-palmetto and saw palmetto habitat. Herbaceous cover in disturbed habitat was diverse and included only small amounts of wiregrass. Recently burned oak-palmetto plots had much higher herbaceous cover than unburned oak-palmetto plots (Table 2).

Many plots in each habitat type had no tortoises, although all but one plot had at least one active or inactive burrow. The plot with no active burrows had active burrows nearby. Standard deviations of densities of tortoises were higher than the means for each habitat type during each season (Table 2). Mean adult and subadult densities of tortoises were usually higher in disturbed habitat than in most oak-palmetto and saw palmetto habitat. Densities of tortoises in saw palmetto habitat exceeded those in oak-palmetto habitat in most seasons and time-since-fire categories, or the differences were small (Table 2). No consistent trends in densities of tortoises occurred among time-since-fire classes in oak-palmetto or saw palmetto habitat.

Only four habitat variables had significant correlations with either fall or spring densities of tortoises and the correlation coefficients were small (Table 3). Only non-*Aristida* cover had significant correlations in both seasons. Time since fire was correlated with herbaceous cover ($r = -0.40$, $P < 0.01$, $N = 103$) and non-*Aristida* herbaceous cover ($r = -0.25$, $P = 0.01$, $N = 103$). Herbaceous cover was highest in plots burned within three years of the study. Herbaceous cover declined more with increasing time-since-fire classes in oak-palmetto habitats than in saw palmetto habitats.

Average densities of hatchling and juvenile burrows were 0.9 (SD = 1.6) burrows/ha for disturbed habitat, 0.7 (SD = 1.3) burrows/ha for oak-palmetto habitat, and 2.3 (SD = 3.6) burrows/ha for saw palmetto habitat. Average densities of hatchling and juvenile burrows were 0.9 (SD = 1.6) burrows/ha in recently burned habitat, 1.0 (SD = 1.7) burrows/ha in habitat burned 4-7 years ago, and 2.5 (SD = 3.7) burrows/ha in habitat unburned for >7 years.

TABLE 3. Spearman rank order correlations (r) between habitat variables and fall and spring gopher tortoise densities on John F. Kennedy Space Center, Florida.

Habitat Parameter	Fall N = 103 r	Spring N = 103 r
% scrub oak cover	-0.21 ($P \leq 0.05$)	-0.11 ($P > 0.05$)
% herbaceous cover	0.06 ($P > 0.05$)	0.23 ($P \leq 0.01$)
% non- <i>Aristida</i> herbaceous cover	0.20 ($P \leq 0.05$)	0.23 ($P \leq 0.01$)
Mild disturbance	0.02 ($P > 0.05$)	0.22 ($P \leq 0.05$)

DISCUSSION

Active gopher tortoise burrows occurred within or adjacent to all study plots suggesting that tortoises were widely distributed in these habitats. Only 20-29% of these burrows were occupied (Breininger et al., 1991a), probably because tortoises used several burrows and frequently relocated to new burrows (e.g., McRae et al., 1981; Douglass, 1990; Diemer, 1992b). Seasonal habitat shifts may have occurred but cannot be determined given the high variation in densities of tortoises. Herbaceous cover had greater influence on density of tortoises elsewhere (Auffenberg and Iverson, 1979; Garner and Landers, 1981) than we observed among our plots. Herbaceous cover within plots may be a poor indicator of available food resources, because swale marshes were near plots, and tortoises fed in them (Breininger et al., 1988).

Densities of tortoises were higher in disturbed habitats than in oak-palmetto and saw palmetto habitats, as expected (Auffenberg and Franz, 1982). Disturbed habitats have fewer shrubs and a more abundant and diverse herbaceous layer than oak-palmetto and saw palmetto habitats (Breininger and Schmalzer, 1990) which may explain the higher densities of tortoises in disturbed habitat. Tortoises will move long distances to find nesting sites, and they select sunny openings (Hallinan, 1923; Landers and Speake, 1980; Landers et al., 1980). The positive correlation between density of tortoises in spring and mild disturbance in oak-palmetto and saw palmetto habitats may be related to nest site selection or basking needs. Open areas, common in disturbed habitats, are rare in most oak-palmetto and saw palmetto habitats except for small areas showing mild soil disturbance associated with historical logging or turpentine activities (Schmalzer and Hinkle, 1992a, b). Disturbed habitat, which had major soil disturbance, is occupied by tortoises for at least 20 yr after clearing; the regenerated habitat often burns poorly and can become less suitable for

tortoises and other species of conservation concern (Breininger and Schmalzer, 1990).

The need for periodic fire to maintain suitable tortoise habitat is well established (Landers, 1980; Landers and Speake, 1980; Auffenberg and Franz, 1982; Diemer, 1986). Densities of tortoises, however, were not always higher in the most recently burned plots, and tortoises occurred in areas unburned for more than 20 years. Food resources were more abundant in recently burned oak-palmetto and saw palmetto plots than in unburned plots, but even unburned plots were near swale marshes where food was abundant.

Densities of tortoises were not usually higher in oak-palmetto (well-drained) than in saw palmetto (poorly-drained) plots in spring or fall. We reported that tortoises did not select well-drained sites in winter (Breininger et al., 1991a). Most studies suggest that well-drained conditions are required by gopher tortoises (e.g., Landers and Speake, 1980; Auffenberg and Franz, 1982), although some note exceptions (Means, 1982; Diemer, 1986). Also, tortoises had slightly higher densities in saw palmetto than in oak-palmetto in another study on KSC, perhaps because more food is near burrows in saw palmetto habitat (Giovanetto, 1988).

Most studies of habitat suitability focus on density comparisons, assuming that density decreases from more suitable to less suitable habitat (Andrewartha and Birch, 1954; Wynne-Edwards, 1962; Flather and Hoekstra, 1985). Measures of reproductive success and survival are important indicators of habitat suitability (Van Horne, 1983). Subadult and adult tortoises were recruited into the study population 5-60 yr before the study when habitat conditions may have been different. Inferences from adult and subadult tortoise densities may not always be indicators of areas suitable for reproductive success. Our results suggested some recruitment of tortoises occurred recently across a range of habitat conditions based on hatchling and juvenile burrows.

Other studies used untested correction factors (for review see Burke and Cox, 1988; Burke, 1989) or reported densities within colonies rather than average densities by habitat; thus, densities are not compared easily across studies. Areas of habitat needed to maintain a population size of 40–50 individuals was estimated to be 10–20 ha (Cox et al., 1987) based on home ranges in sandhill (McRae et al., 1981). Tortoises may require at least 30–35 ha of scrub to support 40–50 adults and subadults, based on average densities at KSC. Additional secondary habitat, such as marshes, may be needed because marshes may provide an important food source.

Acknowledgments.—This paper summarizes a study (Breininger et al., 1988) funded by the Florida Game and Fresh Water Fish Commission (FGFWFC). We thank D. Cook, J. Cox, J. Diemer, and B. Gruver of the FGFWFC for their contributions. We also thank W. Knott III and A. Koller, Jr. of NASA. Field assistance was provided by D. Rydene, B. Blonder, L. Maull, and L. Kulyassa. Helpful comments on the manuscript were provided by R. Smith and D. Cook.

LITERATURE CITED

- ABRAHAMSON, W. G., AND D. C. HARTNETT. 1990. Pine flatwoods and dry prairies. In R. L. Myers and J. J. Ewel (eds.), *Ecosystems of Florida*, pp. 103–149. Univ. Central Florida Press, Orlando.
- ANDREWARTHA, H. G., AND L. C. BIRCH. 1954. *The Distribution and Abundance of Animals*. Univ. Chicago Press, Chicago. 782 pp.
- AUFFENBERG, W., AND J. B. IVERSON. 1979. Demography of terrestrial turtles. In M. Harless and H. Morlock (eds.), *Turtles: Perspectives and Research*, pp. 541–569. Wiley, New York.
- , AND R. FRANZ. 1982. The status and distribution of the gopher tortoise (*Gopherus polyphemus*). In R. B. Bury (ed.), *North American Tortoises: Conservation and Ecology*, pp. 95–126. U.S. Fish Wildl. Serv., Wildl. Res. Rep. 12.
- BREININGER, D. R., AND P. A. SCHMALZER. 1990. Effects of fire and disturbance on vegetation and bird communities in a Florida oak/palmetto scrub. *Amer. Midl. Natur.* 123:64–74.
- , D. A. RYDENE, AND C. R. HINKLE. 1988. Burrow and habitat relationships of the gopher tortoise in coastal scrub and slash pine flatwoods on Merritt Island, Florida. *Florida Game and Fresh Water Fish Comm. Nongame Wildl. Program Final Rep.* 238 pp.
- , AND C. R. HINKLE. 1991a. Estimating occupancy of gopher tortoise (*Gopherus polyphemus*) burrows in coastal scrub and slash pine flatwoods. *J. Herpetol.* 25:317–321.
- , M. PROVANCHIA, AND R. B. SMITH. 1991b. Mapping Florida scrub jay habitat for purposes of land use management. *Photogrammetric Engineering and Remote Sensing* 57:1467–1474.
- BURKE, R. 1989. Burrow-to-tortoise conversion factors: comparison of three gopher tortoise survey techniques. *Herpetol. Rev.* 20:92–94.
- , AND J. COX. 1988. Evaluation and review of field techniques used to study and manage gopher tortoises. In *Management of Reptiles, Amphibians, and Small Mammals in North America*, pp. 205–215. U.S. Dept. Agr., For. Serv. Gen. Tech. Rep. RM-166.
- COX, J., D. INKLEY, AND R. KAUTZ. 1987. Ecology and habitat protection needs of gopher tortoise (*Gopherus polyphemus*) populations found on land slated for large-scale development in Florida. *Florida Game and Fresh Water Fish Commission Nongame Wildlife Program Tech. Rep. No. 4*. 69 pp.
- DIEMER, J. E. 1986. The ecology and management of the gopher tortoise in southeastern United States. *Herpetologica* 42:125–133.
- , 1992a. Demography of the tortoise (*Gopherus polyphemus*) in northern Florida. *J. Herpetol.* 26: 281–289.
- , 1992b. Home range and movements of the tortoise (*Gopherus polyphemus*) in northern Florida. *J. Herpetol.* 26:158–165.
- DOUGLASS, J. F. 1990. Patterns of mate-seeking and aggression in a southern Florida population of the gopher tortoise, *Gopherus polyphemus*. *Proc. Symp. Desert Tortoise Council* 1986:155–199.
- FLATHER, C. H., AND T. W. HOEKSTRA. 1985. Evaluating population-habitat models using ecological theory. *Wildl. Soc. Bull.* 13:121–130.
- GARNER, J. A., AND J. L. LANDERS. 1981. Foods and habitat of the gopher tortoise in southwestern Georgia. *Proc. Ann. Conf. S.E. Assoc. Fish and Wildl. Agen.* 35:120–133.
- GIOVANETTO, L. A. 1988. Habitat use by the gopher tortoise (*Gopherus polyphemus* Daudin) in scrub and slash pine flatwoods on John F. Kennedy Space Center. M.S. Thesis, Fla. Inst. of Technol., Melbourne.
- HALLINAN, T. 1923. Observations made in Duval County, northern Florida, on the gopher tortoise (*Gopherus polyphemus*). *Copeia* 1923:11–20.
- LANDERS, J. L. 1980. Recent research on the gopher tortoise and its implications. In R. Franz and R. J. Bryant (eds.), *The Dilemma of the Gopher Tortoise—Is There a Solution?*, pp. 8–14. *Proc. of First Ann. Mtg., Gopher Tortoise Council*, Auburn Univ., Alabama.
- , AND D. W. SPEAKE. 1980. Management needs of sandhill reptiles in southern Georgia. *Proc. Ann. Conf. S.E. Assoc. Fish and Wildl. Agen.* 34:515–529.
- , J. H. GARNER, AND W. A. McRAE. 1980. Reproduction of gopher tortoise (*Gopherus polyphemus*) in southwestern Georgia. *Herpetologica* 36: 353–361.
- MACDONALD, L. A., AND H. R. MUSHINSKY. 1988. Foraging ecology of the gopher tortoise, *Gopherus polyphemus*, in a sandhill habitat. *Herpetologica* 44: 345–353.
- McRAE, W. A., J. L. LANDERS, AND J. A. GARNER. 1981. Movement patterns and home range of the gopher tortoise. *Amer. Midl. Natur.* 106:165–179.
- MEANS, D. B. 1982. Responses to winter burrow flooding of the gopher tortoise (*Gopherus polyphemus* Daudin). *Herpetologica* 38:521–525.

- MUELLER-DOMDOIS, D., AND H. ELLENBERG. 1974. *Aims and Methods of Vegetation Ecology*. Wiley, New York. 547 pp.
- MYERS, R. L. 1990. Scrub and high pine. In R. L. Myers and J. J. Ewel (eds.), *Ecosystems of Florida*, pp. 150-193. Univ. Central Florida Press, Orlando.
- SCHMALZER, P. A., AND C. R. HINKLE. 1992a. Recovery of oak-saw palmetto scrub after fire. *Castanea* 57: 158-173.
- _____, AND _____. 1992b. Species composition and structure of oak-saw palmetto scrub vegetation. *Castanea* 57:220-251.
- SPSS INC. 1988. *SPSS/PC+ V2.0 Base Manual for the IBM PC/XT/AT and PS/2*. SPSS Inc., Chicago, Illinois.
- VAN HORNE, B. 1983. Density as a misleading indicator of habitat quality. *J. Wildl. Manage.* 47:813-901.
- WYNNE-EDWARDS, V. C. 1962. *Animal Dispersion in Relation to Social Behavior*. Oliver and Boyd, Edinburgh, Scotland. 653 pp.

Accepted: 14 October 1993.